

## Grant H Garcia, MD

2409 North 45<sup>th</sup> Street,  
Seattle, WA 98103  
Wallingford: 206-633-8100  
Ballard: 206-784-8833



### POSTOPERATIVE INSTRUCTIONS

#### **CAPSULAR RELEASE AND MANIPULATION UNDER ANESTHESIA**

##### **DIET**

- Begin with clear liquids and light foods (jello, soups, etc.)
- Progress to your normal diet if you are not nauseated

##### **WOUND CARE**

- Maintain your operative dressing, loosen bandage if swelling of the foot or ankle occurs
- It is normal for the knee to bleed and swell following surgery – if blood soaks onto the bandage, do not become alarmed – reinforce with additional dressing
- Remove surgical dressing on the second post-operative day – if minimal drainage is present, apply waterproof band-aids over incisions and change daily
- To avoid infection, keep surgical incisions clean and dry – you may shower by placing waterproof band-aids over incision areas. Please remember to change band-aids daily
- NO immersion of operative leg (i.e. bath) \*Brace may come off to shower

##### **MEDICATIONS**

###### **\*Do not drive a car or operate machinery while taking the narcotic medication\***

- Pain medication is injected into the wound and knee joint during surgery – this will wear off within 8-12 hours
- Most patients will require some narcotic pain medication for a short period of time
- Primary Medication = Norco (Hydrocodone)
  - Take 1 – 2 tablets every 4 – 6 hours as needed
  - Max of 12 pills per day
  - Plan on using it for 2 to 5 days, depending on level of pain
  - Do NOT take additional Tylenol (Acetaminophen) while taking Norco or Vicodin
- Common side effects of the pain medication are nausea, drowsiness, and constipation – to decrease the side effects, take medication with food.
- If constipation occurs, consider taking an over-the-counter laxative such as prune juice, Senekot, Colace (or Periocolase), or Miralax
- If you are having problems with nausea and vomiting, contact the office to possibly have your medication changed
  - For nausea, take prescribed Zofran / Phenergan
- Ibuprofen 600-800mg (i.e., Advil) may be taken in between the narcotic pain medication to help smooth out the postoperative “peaks and valleys”, reduce overall amount of pain medication required, and increase the time intervals between narcotic pain medication usage

##### **Activity**

- Elevate the operative leg to chest level whenever possible to decrease swelling
- Place pillows under knees (i.e., do not maintain knee in a flexed or bent position), but rather place pillows under foot/ankle to elevate leg

- Use crutches to assist with walking as needed
- Do not engage in activities which increase knee pain/swelling (prolonged periods of standing or walking) over the first 7-10 days following surgery
- Avoid long periods of sitting (without leg elevated) or long distance traveling for 2 weeks
- May return to sedentary work or school 3-4 days after surgery, if pain is tolerable

### **Ice Therapy**

- Begin immediately after surgery  
Use Duracold ice packs (if machine not prescribed) every 2 hours for 20 minutes daily until your first post-operative visit – remember to keep leg elevated to level of chest while icing

### **Exercise**

- Begin exercises 24 hours after surgery (straight leg raises, quad sets, heel slides and ankle pumps) unless otherwise instructed
- Discomfort and knee stiffness is normal for a few days following surgery – it is safe and, in fact preferable to bend your knee (unless otherwise instructed by physician)
- Complete exercises 3-4 times daily until your first postoperative visit – your motion goals are to have complete extension (straightening) and 90 degrees of flexion (bending) at your first postoperative visit unless otherwise instructed
- Do ankle pumps continuously throughout the day to reduce the possibility of a blood clot in your calf (extremely uncommon)
- ***Formal physical therapy (PT) will begin first day after surgery 5 times per week for 2-3 weeks.***

## **CONTACTING OUR OFFICE**

### **EMERGENCIES\*\***

\*Contact Dr. Garcia at 206-633-8100 if any of the following are present:

- Painful swelling or numbness (note that some swelling and numbness is normal)
- Unrelenting pain
- Fever (over 101° - it is normal to have a low grade fever for the first day or two following surgery) or chills
- Redness around incisions
- Color change in foot or ankle
- Continuous drainage or bleeding from incision (a small amount of drainage is expected)
- Difficulty breathing
- Excessive nausea/vomiting
- Calf pain

\*\*If you have an emergency after office hours or on the weekend, contact the same office number (206-633-8100) and you will be connected to our page service – this will contact Dr. Garcia or one of his physician assistants. If he is unavailable. Do NOT call the hospital or surgicenter.

\*\*If you have an emergency that requires immediate attention, proceed to the nearest emergency room.

### **FOLLOW-UP CARE/QUESTIONS**

\*If you do not already have a post-operative appointment scheduled, please contact the office immediately at 206-633-8100 to speak our Administrative Assistant or visit our website [www.grantgarciamd.com](http://www.grantgarciamd.com)

\*Typically post-operative appointments are made for 10-14 days following the date of surgery

\*If you have any further questions, contact the office directly.