

Grant H Garcia, MD
 2409 North 45th Street,
 Seattle, WA 98103
 Wallingford: 206-633-8100
 Ballard: 206-784-8833



ACL PATELLAR TENDON AUTOGRAFT RECONSTRUCTION REHABILITATION PROTOCOL

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I 0-4 weeks	Full in Brace	0-1 week: Locked in full extension for ambulation and sleeping 1-4 weeks: Unlocked for ambulation, remove for sleeping**	As tolerated	Heel slides, quad/hamstring sets, patellar mobs, gastroc/soleus stretch SLR w/ brace in full extension until quad strength prevents extension lag Side-lying hip/core
PHASE II 4-12 weeks	Full	Discontinue at day 28 if patient has no extension lag	Full	Begin toe raises, closed chain quads, balance exercises, hamstring curls, stationary bike, step-ups, front and side planks; advance hip/core
PHASE III 12-16 weeks	Full	None	Full	Advance closed chain strengthening Progress proprioception activities Begin stairmaster, elliptical at 8 wks; Running straight ahead at 12 weeks
PHASE IV 16-24 weeks	Full	None	Full	16 wks: Begin jumping 20 wks: Advance to sprinting, backward running, cutting/pivoting/changing direction, initiate plyometric program and sport-specific drills 22 wks: Advance as tolerated FSA completed at 22 wks***
PHASE V > 6 months	Full	None	Full and pain-free	Gradual return to sports participation after completion of FSA Maintenance program based on FSA

*Modified with concomitantly performed meniscus repair/transplantation or articular cartilage procedure

**Brace may be removed for sleeping after first post-operative visit (day 7-10)

***Completion of FSA (Functional Sports Assessment) not mandatory, but recommended at approx 22 wks post-op for competitive athletes returning to play after rehab