Grant H Garcia, MD

2409 North 45th Street, Seattle, WA 98103

Wallingford: 206-633-8100 Ballard: 206-784-8833



ACL HAMSTRINGS & QUAD TENDON AUTOGRAFT RECONSTRUCTION REHABILITATION PROTOCOL

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I 0-4 weeks	As tolerated with crutches*	 0-1 week: Locked in full extension for ambulation and sleeping 1-4 weeks: Unlocked for ambulation, remove for sleeping** 	As tolerated	Heel slides, quad/hamstring sets, patellar mobs, gastroc/soleus stretch SLR w/ brace in full extension until quad strength prevents extension lag
PHASE II 4-12 weeks	Full, progressing to normal gait pattern	Discontinue at day 28 if patient has no extension lag	Main full extension and progressive flexion	Progress Phase I Begin toe raises, closed chain quads, balance exercises, hamstring curls, stationary bike, step-ups, front and side planks, hip/glute/core, pool
PHASE III 12-16 weeks	Full, without use of crutches and with a normalized gait pattern	None	Gain full and pain-free	Advance closed chain strengthening, progress proprioception activities Begin stairmaster, elliptical and running straight ahead
PHASE IV 16-24 weeks	Full	None	Full and pain- free	16 wks: Begin jumping 20 wks: Advance running to sprinting, backward running, cutting/pivoting/changing direction, initiate plyometric program and sport-specific drills 22 wks: Advance as tolerated FSA completed at 22 wks***
PHASE V > 6 months	Full	None	Full and pain- free	Gradual return to sports participation after completion of FSA Maintenance program based on FSA

^{*}Modified with concomitantly performed meniscus repair/transplantation or articular cartilage procedure

^{**}Brace may be removed for sleeping after first post-operative visit (day 7-10)

^{***}Completion of FSA (Functional Sports Assessment) not mandatory, but recommended at approx 22 wks post-op for competitive athletes returning to play after rehab