

**Grant H Garcia, MD**  
 2409 North 45<sup>th</sup> Street,  
 Seattle, WA 98103  
 Wallingford: 206-633-8100  
 Ballard: 206-784-8833



## **DISTAL CLAVICLE OPEN REDUCTION INTERNAL FIXATION PT**

**\*\*Distal clavicle fractures are less secure than mid-shaft clavicle fractures and require a more conservative protocol**

	<b>RANGE OF MOTION</b>	<b>IMMOBILIZER</b>	<b>EXERCISES</b>
<b>PHASE I</b> 0-6 weeks	Gentle supine PROM as tolerated avoiding horizontal adduction	<b>0-6 weeks:</b> Remove only for exercise and hygiene which should be performed supine with gravity eliminated	Closed chain scapular stabilizers, deltoid and rotator cuff while supine or with gravity eliminated  ROM and strengthening: elbow and shoulder isometric exercises
<b>PHASE II</b> 6-14 weeks	AROM as tolerated in prone position	None	Progress Phase I exercises  Active assisted strength work in all ROMs  Begin vertical positioned strengthening at 10 weeks
<b>PHASE III</b> 14-24 weeks	Progress to full AROM in all planes	None	Advance strengthening as tolerated  Progress Phase II and Phase I exercises  Begin sport-specific exercises after week 20 when cleared by MD