

Autologous Cultured Chondrocyte Implantation (MACI®) Patella and/or Trochlea Rehab Protocol

Description of Procedure: Two stage technique. **Stage 1:** A small amount of the patient's own articular cartilage is harvested and through cell culturing techniques, the cell number is increased from a few hundred thousand to over 10 million cells. **Stage 2:** These cultured chondrocytes are then re-implanted in the knee on a porcine collagen membrane patch placed within the defect. The cells then gradually form hyaline-like cartilage to resurface areas of prior cartilage loss.

Safety Warning: Do not overload the implant, especially with shear forces, as patch delamination could occur. Early approved ROM is important to avoid excessive scarring and to stimulate the chondrocytes to form hyaline-like extracellular matrix. If the patient has had concomitant tibial tuberosity surgery, the weight bearing restrictions of that procedure take precedence.

	Weight Bearing	Brace	ROM	Therapeutic Exercise
Phase I: 0 to 6 Weeks	Weight bearing as tolerated once a straight leg raise can be performed without extension lag with the use of two crutches <i>**Avoid stairs and inclines for 6 months post-operatively</i>	Brace is worn until whenever ambulating for 6 weeks post-operatively	0 to 3 Weeks: CPM: use for 8 hours per day at 1 cycle/minute - begin at 0 to 30° increasing as rapidly as possible to re-establish full motion anytime patient does not feel 'stretch' <i>**Goal: To achieve active range of motion as soon as tolerated</i>	Prone hangs, heel props, quad sets, SLR, hamstring isometrics - complete exercises in brace if quad control is inadequate; core proximal program; normalize gait; FES biofeedback as needed <i>** Incorporate use of stationary bike (high seat, low resistance) and patellar mobilization exercises after surgical dressing is removed</i>
Phase II: 6 to 12 Weeks	Progression to full weight bearing with normalized gait pattern; no limping (avoid stairs and inclines for 6 months)	None	Full active range of motion	Progress bilateral closed chain strengthening using resistance less than patient's body weight, progress to supine unilateral leg press with low weight, high reps; continue opened chain knee strengthening <i>**NO squats, wall slides, lunges or knee extension exercises</i>
Phase III: 12 Weeks to 12 Months	Full, with a normalized gait pattern (avoid stairs and inclines for 6 months)	None	Full active range of motion	12 Weeks to 6 Months: Advance bilateral and unilateral closed chain exercises with emphasis on concentric/eccentric control, continue with biking, elliptical, and walking on treadmill, progress

	<i>**May progress stairs if adequate quad strength no pain or swelling</i>			balance activities 6 to 12 Months: Advance strength training <i>**May progress stairs if adequate quad strength no pain or swelling</i>
Phase IV: 12 to 24 Months	Full, with a normalized gait pattern	None	Full active range of motion	Continue Strength Training: Initiate light jogging - start with 2 minute walk/2 minute jog, emphasize sport-specific training; emphasize single leg loading, plyometrics, begin agility pro- gram - <u>high impact activities (basketball, tennis, etc.) may begin at 12 to 24 months after passing a pain free functional progression test</u>

Progression back to sport is dependent on case per case basis and determined by Dr. Garcia. If pain or swelling occurs patient is expected to stop causative activity and follow-up with our office.

Leg extension exercises with resistance are not allowed indefinitely.