

Grant H Garcia, MD
2409 North 45th Street,
Seattle, WA 98103
Wallingford: 206-633-8100
Ballard: 206-784-8833



Post-Operative Rehabilitation Guidelines for Massive Rotator Cuff Tears & Superior Capsular Reconstruction

- 1-6 Weeks: Sling Immobilization
Active ROM Elbow, Wrist and Hand
True Passive (ONLY) ROM Shoulder
 Pendulums,
 Supine Elevation in Scapular plane = 140 degrees
 External Rotation = 40 degrees
Scapular Stabilization exercises (side-lying)
Deltoid isometrics in neutral (submaximal) as ROM improves
No Pulley/Canes until 6 weeks post-op
- 6-12 Weeks: Discontinue Sling
Active Assist to Active ROM Shoulder As Tolerated
 Elevation in scapular plane and external rotation to tolerance
 Begin internal rotation as tolerated
 Light stretching at end ranges
Cuff Isometrics with the arm at the side
Upper Body Ergometer
- 3-12 Months Advance to full ROM as tolerated with passive stretching at end ranges
Advance strengthening as tolerated: isometrics → bands → light weights (1-5
 lbs); 8-12 reps/2-3 sets per rotator cuff, deltoid, and scapular stabilizers
Only do strengthening 3x/week to avoid rotator cuff tendonitis
Begin eccentrically resisted motions, plyometrics (ex. Weighted ball toss),
proprioception (es. body blade)
Begin sports related rehab at 4 ½ months, including advanced conditioning
Return to throwing at 6 months
Throw from pitcher's mound at 9 months
Collision sports at 9 months
MMI is usually at 12 months post-op